

# Introduction to Global Warming and Carbon Footprinting

Dr. Richard Venditti

Forest Biomaterials

North Carolina State University

Source if no other reference appears:

**This Summary for Policymakers should be cited as:**

IPCC, 2007: Summary for Policymakers. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.]

# Global Warming Changes





Figure SPM.3. Observed changes in (a) global average surface temperature, (b) global average sea level from tide gauge (blue) and satellite (red) data and (c) Northern Hemisphere snow cover for March–April. All changes are relative to corresponding averages for the period 1961–1990. Smoothed curves represent decadal average values while circles show yearly values. The shaded areas are the uncertainty intervals estimated from a comprehensive analysis of known uncertainties (a and b) and from the time series (c). (FAQ 3.1, Figure 1, Figure 4.2, Figure 5.13)

# Global Carbon Cycle


- ***The **velocity** of climate change may have more impact than the absolute value of the changes***

# Changing temperature in California.



SR Loarie *et al. Nature* **462**, 1052-1055 (2009) doi:10.1038/nature08649


# Global Warming Predictions



**Figure SPM.5.** Solid lines are multi-model global averages of surface warming (relative to 1980–1999) for the scenarios A2, A1B and B1, shown as continuations of the 20th century simulations. Shading denotes the  $\pm 1$  standard deviation range of individual model annual averages. The orange line is for the experiment where concentrations were held constant at year 2000 values. The grey bars at right indicate the best estimate (solid line within each bar) and the likely range assessed for the six SRES marker scenarios. The assessment of the best estimate and likely ranges in the grey bars includes the AOGCMs in the left part of the figure, as well as results from a hierarchy of independent models and observational constraints. (Figures 10.4 and 10.29)

# Changes in GHGs

- Global atmospheric concentrations of carbon dioxide, methane and nitrous oxide have increased markedly as a result of human activities since 1750
- Now far exceed pre-industrial values determined from ice cores spanning many thousands of years
- The global increases in
  - carbon dioxide concentration are due primarily to fossil fuel use and land use change,
  - Methane and nitrous oxide are primarily due to agriculture.



*Figure SPM.1. Atmospheric concentrations of carbon dioxide, methane and nitrous oxide over the last 10,000 years (large panels) and since 1750 (inset panels). Measurements are shown from ice cores (symbols with different colours for different studies) and atmospheric samples (red lines). The corresponding radiative forcings are shown on the right hand axes of the large panels. [Figure 6.4]*

# Global Carbon Cycle





Fig. 1 Estimates of the global pools and fluxes between them.<sup>1,4,5,7,152</sup>

# Global Warming



**Figure 2.** Ice core record from Vostok, Antarctica, showing the near-simultaneous rise and fall of Antarctic temperature and CO<sub>2</sub> levels through the last 350,00 years, spanning three ice age cycles. However, there is a lag of several centuries between the time the temperature increases and when the CO<sub>2</sub> starts to increase. Image credit: [Siegenthaler et al., 2005, Science](#)



# Global Warming Potential (GWP)

- relative measure of how much heat a [greenhouse gas](#) traps in the atmosphere.
- compares the amount of heat trapped by a certain mass of the [gas](#) in question to the amount of heat trapped by a similar mass of [carbon dioxide](#).
- commonly determined over a span of 20, 100 or 500 years.
- GWP is expressed as a factor of carbon dioxide (whose GWP is standardized to 1).

# Radiative Forcing Capacity (RF) and GWP

- RF = the amount of energy per unit area, per unit time, absorbed by the greenhouse gas, that would otherwise be lost to space
- GWP is the ratio of the time-integrated radiative forcing from the instantaneous release of 1 kg of a trace substance relative to that of 1 kg of a reference gas

$$GWP_i \equiv \frac{\int_0^{TH} RF_i(t) dt}{\int_0^{TH} RF_r(t) dt} = \frac{\int_0^{TH} a_i \cdot [C_i(t)] dt}{\int_0^{TH} a_r \cdot [C_r(t)] dt}$$

- where  $TH$  is the time horizon,
- $RF_i$  is the global mean RF of component  $i$ ,
- $a_i$  is the RF per unit mass increase in atmospheric abundance of component  $i$  (radiative efficiency),
- $[C_i(t)]$  is the time-dependent abundance of  $i$ ,
- and the corresponding quantities for the reference gas ( $r$ ) in the denominator.


# Global Warming Potential Values

**Table 2.14.** Lifetimes, radiative efficiencies and direct (except for CH<sub>4</sub>) GWP's relative to CO<sub>2</sub>. For ozone-depleting substances and their replacements, data are taken from IPCC/TEAP (2005) unless otherwise indicated.

Industrial Designation or Common Name (years)	Chemical Formula	Lifetime (years)	Radiative Efficiency (W m <sup>-2</sup> ppb <sup>-1</sup> )	Global Warming Potential for Given Time Horizon			
				SAR <sup>a</sup> (100-yr)	20-yr	100-yr	500-yr
Carbon dioxide	CO <sub>2</sub>	See below <sup>a</sup>	1.4x10 <sup>-5</sup>	1	1	1	1
Methane <sup>c</sup>	CH <sub>4</sub>	12 <sup>c</sup>	3.7x10 <sup>-4</sup>	21	72	25	7.6
Nitrous oxide	N <sub>2</sub> O	114	3.03x10 <sup>-3</sup>	310	289	298	153
<b>Substances controlled by the Montreal Protocol</b>							
CFC-11	CCl <sub>3</sub> F	45	0.25	3,800	6,730	4,750	1,620
CFC-12	CCl <sub>2</sub> F <sub>2</sub>	100	0.32	8,100	11,000	10,900	5,200
CFC-13	CClF <sub>3</sub>	640	0.25		10,800	14,400	16,400
CFC-113	CCl <sub>2</sub> FCClF <sub>2</sub>	85	0.3	4,800	6,540	6,130	2,700
CFC-114	CClF <sub>2</sub> CClF <sub>2</sub>	300	0.31		8,040	10,000	8,730
CFC-115	CClF <sub>2</sub> CF <sub>3</sub>	1,700	0.18		5,310	7,370	9,990
Halon-1301	CBrF <sub>3</sub>	65	0.32	5,400	8,480	7,140	2,760
Halon-1211	CBrClF <sub>2</sub>	16	0.3		4,750	1,890	575
Halon-2402	CBrF <sub>2</sub> CBrF <sub>2</sub>	20	0.33		3,680	1,640	503
Carbon tetrachloride	CCl <sub>4</sub>	26	0.13	1,400	2,700	1,400	435
Methyl bromide	CH <sub>3</sub> Br	0.7	0.01		17	5	1
Methyl chloroform	CH <sub>3</sub> CCl <sub>3</sub>	5	0.06		506	146	45
HCFC-22	CHClF <sub>2</sub>	12	0.2	1,500	5,160	1,810	549
HCFC-123	CHCl <sub>2</sub> CF <sub>3</sub>	1.3	0.14	90	273	77	24
HCFC-124	CHClF <sub>2</sub> CF <sub>3</sub>	5.8	0.22	470	2,070	609	185
HCFC-141b	CH <sub>3</sub> CCl <sub>2</sub> F	9.3	0.14		2,250	725	220
HCFC-142b	CH <sub>3</sub> CClF <sub>2</sub>	17.9	0.2	1,800	5,490	2,310	705
HCFC-225ca	CHCl <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>	1.9	0.2		429	122	37
HCFC-225cb	CHClF <sub>2</sub> CClF <sub>2</sub>	5.8	0.32		2,030	595	181
<b>Hydrofluorocarbons</b>							
HFC-23	CHF <sub>3</sub>	270	0.19	11,700	12,000	14,800	12,200
HFC-32	CH <sub>2</sub> F <sub>2</sub>	4.9	0.11	650	2,330	675	205
HFC-125	CHF <sub>2</sub> CF <sub>3</sub>	29	0.22	2,800	6,350	3,500	1,100

# Radiative Forcing

- rate of energy change per unit area of the globe as measured at the top of the atmosphere
- expressed in units of Watts per square metre



**Figure SPM.2.** Global average radiative forcing (RF) estimates and ranges in 2005 for anthropogenic carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and other important agents and mechanisms, together with the typical geographical extent (spatial scale) of the forcing and the assessed level of scientific understanding (LOSU). The net anthropogenic radiative forcing and its range are also shown. These require summing asymmetric uncertainty estimates from the component terms, and cannot be obtained by simple addition. Additional forcing factors not included here are considered to have a very low LOSU. Volcanic aerosols contribute an additional natural forcing but are not included in this figure due to their episodic nature. The range for linear contrails does not include other possible effects of aviation on cloudiness. (2.9, Figure 2.20)

# Carbon Footprint: Impact Assessment Method

- **Partial life cycle analysis**
- **Historically:** the total set of greenhouse gas (GHG) emissions caused by an organization, event, product or person (UK Carbon Trust, 2009)
- **Practically:** A measure of the total amount of carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) emissions of a defined population, system or activity, considering all relevant sources, sinks and storage within the spatial and temporal boundary of the population, system or activity of interest. Calculated as carbon dioxide equivalent (CO<sub>2</sub>e) using the relevant 100-year global warming potential (GWP100) (Wright et al, Carbon Mgmt, 2011)




# Carbon Footprint: Impact Assessment Method

- **IPCC is the leading authority in evaluating the science behind GWP**

Revision Year	CO <sub>2</sub> equivalents for CH <sub>4</sub>	CO <sub>2</sub> equivalents for N <sub>2</sub> O
1996	21	310
2001	23	296
2006	25	298



# Carbon Footprint: A Material Balance of GHG's




Carbon footprint = Emissions- Absorption (kg CO2 equivalents)



# Carbon Footprint: Impact Assessment Method

- Typically, a carbon footprint does not consider biogenic (from living processes) carbon nor does it consider CO<sub>2</sub> emissions from the burning or decay of the biogenic material (they balance each other)
- Biogenic material decay/burning that produces methane or N<sub>2</sub>O must be considered









# Carbon Footprint: Impact Assessment Method

- **Non renewable resources (coal, oil) are considered since they have been formed over very long time scales and are not being formed over time scales of interest**
- **Materials, transportation, energy often have associated with them carbon emissions**
- **Long term storage of carbon away from the atmosphere is considered a negative C footprint contribution**
- **When one product with a lower C footprint replaces another with larger C footprint, an avoided C input to the atmosphere is claimed, a negative C footprint contribution**



# Carbon Footprint: CO2 list.org

This list may confirm what you know, or may surprise you.  
CO<sub>2</sub> is not caused by others, it is caused by our choices: Heating & cooling; Buying products; Red meat versus chicken and grain; Cars and


<a href="http://CO2LIST.ORG">CO2LIST.ORG</a> Home  <b>Bold shows some interesting items</b>	<b>KILOS OF CO<sub>2</sub></b>  (includes effect of other greenhouse gases)	<b>POUNDS OF CO<sub>2</sub></b>	<b>UNITS OF MEASURE FOR EACH ITEM</b>  (We and most others measure CO <sub>2</sub> by weight. Its size varies, so it can't be measured in volume. For other items we pick appropriate units, shown below.)	What is the answer? Solutions are discussed at <a href="http://Your.CO2List.org">http://Your.CO2List.org</a> Complete sources and calculations are at <a href="http://sfs.CO2List.org">sfs.CO2List.org</a> Data from US, except when the following symbols appear: † Data are from UK ‡ Data are from Australia France has data (in English) for many items, not yet incorporated here. Contact us ... 16 March 2012
--	---	---------------------------------	--	--

## CO<sub>2</sub> POUNDS RELEASED WHILE MAKING PRODUCTS

1 - FOOD	KILOS OF CO <sub>2</sub>	POUNDS OF CO <sub>2</sub>		Pounds of CO <sub>2</sub> per 500 Calories (this is 1/4 of a daily 2,000-Calorie diet)	
<b>Red meat</b>	<b>22</b>	<b>22</b>	Pounds CO <sub>2</sub> per pound of product, or Kilos of CO <sub>2</sub> per kilo of product	12	92% from production of animals & their feed, including N <sub>2</sub> O & methane. Remainder is transport of inputs & meat, and selling. (interesting article by former Texas Ag Commissioner <a href="http://jimhightower.com/node/6901">http://jimhightower.com/node/6901</a> )
<b>Chicken, fish, eggs</b>	<b>6</b>	<b>6</b>		4	81% from production of feed & meat
<b>Dairy</b>	<b>4</b>	<b>4</b>		6	91% from production of feed & animals
<b>Cereals, carbohydrates</b>	<b>3</b>	<b>3</b>		1.5	75% from production of crops
<b>Fruit, vegetables</b>	<b>2</b>	<b>2</b>		4	74% from production of crops
Oils, sweets, condiments	<b>2</b>	<b>2</b>		0.5	74% from production of crops
Balanced Diet				1.7	USDA Food Guide: 53% carbohydrate, 29% oils, 18% protein (here protein is chicken, fish, eggs) <a href="http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2005/2005">http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2005/2005</a>
Source: Weber & Matthews 2008 "Food-Miles and the Relative Climate Farm products (food, cloth, leather, biofuels) release greenhouse gases from and Crutzen et al. 2008 "N <sub>2</sub> O Release...")			<a href="http://pubs.acs.org/doi/full/10.1021/es702969f">http://pubs.acs.org/doi/full/10.1021/es702969f</a> <a href="http://www.iea.org/textbase/nppdf/free/2004/biofuels2004.pdf">http://www.iea.org/textbase/nppdf/free/2004/biofuels2004.pdf</a> <a href="http://www.atmos-chem-phys.net/8/389/2008/acp-8-389-2008.html">http://www.atmos-chem-phys.net/8/389/2008/acp-8-389-2008.html</a>		
(d) methane (CH <sub>4</sub> ) created in animal stomachs and intestines, (e) 2009 report on Biofuels, particularly chapters 6 on land use and 5 on			<a href="http://icsu.org/">http://icsu.org/</a> <a href="http://cip.cornell.edu/biofuels/">http://cip.cornell.edu/biofuels/</a>		
Potato chips‡	2	<b>2</b>	pounds CO <sub>2</sub> per pound of product  kilos CO <sub>2</sub> per kilo of product	Mostly from growing crops: N <sub>2</sub> O from nitrogen-fixing bacteria, fuel  The figures in the section above are larger, and come from a much more complete methodology.	
Orange juice	0.9-1.4	<b>0.9-1.4</b>			
Bottled smoothie‡	1.1	<b>1.1</b>			
Organic new potato‡	0.29	<b>0.29</b>			
Potato, not organic‡	0.24	<b>0.24</b>			
Sources: Carbon Trust, a UK nonprofit, has a summary and Report CTC744.  Orange juice is from a Pepsico study reported in the <i>NY Times</i> .			<a href="http://www.carbon-label.com/individuals/product.html">http://www.carbon-label.com/individuals/product.html</a> <a href="http://www.carbontrust.co.uk/Publicsites/cScape.CT.PublicationsOrd?id=CTC744">http://www.carbontrust.co.uk/Publicsites/cScape.CT.PublicationsOrd?id=CTC744</a> <a href="http://www.nytimes.com/2009/01/22/business/22pepsi.html/">http://www.nytimes.com/2009/01/22/business/22pepsi.html/</a>		

# Carbon Footprint:

Japan's Central Research Institute of the Electric Power Industry's




## **Carbon Footprint Example: Coated Paper (Catalog)**

**Data and Graphs from NCASI LCA P&W Grades, 2010**  
**Software used from NCASI, FEFPRO**

NCASI LCA NA P&W Grades, 2010

# Life Cycle Analysis of Paper: Carbon Footprint Results




# Basic Steps of the Carbon Footprint

- Define the footprint boundary
- Define the scope
- Define the Basis of Calculation
- Begin to complete the Life Cycle Inventory
  - Forest Carbon Changes
  - Wood and Fiber
  - Fuels from Mfg
  - Other Materials
  - Electricity and Steam
  - Mfg waste
  - Product Transport
  - End of Life
- Evaluate Results, Interpret, Report

# Define the footprint boundary

- Cradle to Grave of catalog paper, coated free sheet
- 100 years

**LEGEND:**



**11. AVOIDED EMISSIONS AND OTHERS**

Benefits from sold electricity

Avoided emissions: landfill methane capture and burning for energy

Avoided emissions: product burning for energy

Manufacturing biomass CO<sub>2</sub> emissions

# Define the scope

- 100 years
- Scope 1: all direct GHG emissions from owned production;
- Scope 2: indirect GHG emissions from consumption of purchased electricity, heat or steam; and
- Scope 3: indirect GHG emission from systems such as extraction and production of purchased materials and fuels, transportation in *non-owned* vehicles, or production facilities operated by parties other than the user.



# Define the Basis of Calculation

- Basis of Calculation (BoC) is the metric upon which all of the data input, calculations, and result output are based. For example, a BoC of 1000 kg of product (one metric tonne) means that data input such as quantity of raw material consumed is entered per 1000 kg of production (e.g., a BoC of 1000 kg and log input of 2000 kg means that 2 tonnes of logs are consumed in the production of 1 tonne of product).

<b>Name of this footprint</b>	Coated Freesheet NCASI Number of Uses Cradle to Grave
Product Name	Coated Freesheet
Product Type	coated woodfree
Footprint Boundaries	Cradle to Grave only in this version
Description of a single product	1 mdst (5% water)
Basis of Calculation (BoC)	1 machine-dry short ton (5% water)
Basis of calculation (BoC) expressed as mass (dry kg)	861.82556

# Life Cycle Inventory: Forest Carbon Changes

- Must understand if the land that is being used to provide the amount of wood needed to make paper is being changed such that the net carbon stock on the land for 100 years is changing over many harvests
- Not commonly known, but can be important
- In developed countries, significant proportion is harvested sustainably and many certified

Forest name	Method of determining carbon stock changes	Change in carbon stocks (kg carbon/BoC)	
		Default value	Selected value
<i>Generic Forest</i>	<i>Constant Stock</i>	0	0.00

# Life Cycle Inventory: Wood and Fiber

- Wood and Fiber inputs into manufacturing
- Northern Hardwood chips example, but most cases have multiple inputs

		Proposed defaults	From owned operations	From non-owned operations
Quantity (kg/BoC, dry basis)		<i>No default</i>		261
Moisture content as received (fraction between zero and one)		<i>No default</i>		0.5
Emissions for this fiber source (kg CO <sub>2</sub> eq./kg dry)	Scope 1	<i>0.130</i>		0
	Scope 2	<i>0.090</i>		0
	Scope 3	<i>0.010</i>		0.230
Wet tonnes			0	0.522
Total shipped tonnes			0.522	

	kg CO <sub>2</sub> eq./BoC		
	Scope 1	Scope 2	Scope 3
Emissions for this fiber source	0	0	50.028364
Truck, owned	0		0
Truck, non-owned			4.394349
Rail, owned	0		0
Rail, non-owned			0.1841231
Water inland, owned	0		0
Water inland, non-owned			0
Water ocean, owned	0		0
Water ocean, non-owned			0
<b>Total</b>	<b>0</b>	<b>0</b>	<b>54.606836</b>
<b>Transport only</b>	<b>0</b>		<b>4.5784721</b>

# Life Cycle Inventory: Fuels Consumed

- Coal, example

		Proposed defaults	Burned in owned operations	Burned in non-owned operations
Quantity (GJ HHV/BoC, dry basis)		<i>No default</i>	5.56	
Moisture content as received (fraction between zero and one)		0.1	0.1	
Emissions for this fuel (kg CO <sub>2</sub> eq./GJ HHV)	Combustion	90.32	90.32	N/A
	Pre-combustion	5.382	5.382	N/A
	Total	95.702	95.702	
Transported tons			0.196744515	0
Total transported tons			0.196744515	

	kg CO <sub>2</sub> eq./BoC		
	Scope 1	Scope 2	Scope 3
Fuel-related emissions	425.8628		25.376366
Truck, owned	0		0
Truck, non-owned			0.0453493
Rail, owned	0		0
Rail, non-owned			2.3267772
Water inland, owned	0		0
Water inland, non-owned			0.0545282
Water ocean, owned	0		0
Water ocean, non-owned			0
<b>Total</b>	<b>425.8628</b>		<b>27.80302</b>
<b>Transport only</b>	<b>0</b>		<b>2.4266546</b>

# Life Cycle Inventory: Fuels Consumed

- Black liquor, organic material byproduct of making paper

		Proposed defaults	Burned in owned operations	Burned in non-owned operations
Quantity (GJ HHV/BoC, dry basis)		<i>No default</i>	9.1	
Moisture content as received (fraction between zero and one)		0.35	0.35	
Emissions for this fuel (kg CO <sub>2</sub> eq./GJ HHV)	Combustion	0.637	0.637	N/A
	Pre-combustion	0	0	N/A
	Total	0.637	0.637	
Transported tons			1	0
Total transported tons			1	

	kg CO <sub>2</sub> eq./BoC		
	Scope 1	Scope 2	Scope 3
Fuel-related emissions	4.830908		0
Truck, owned	0		0
Truck, non-owned			0
Rail, owned	0		0
Rail, non-owned			0
Water inland, owned	0		0
Water inland, non-owned			0
Water ocean, owned	0		0
Water ocean, non-owned			0
<b>Total</b>	<b>4.830908</b>		<b>0</b>
<b>Transport only</b>	<b>0</b>		<b>0</b>

# Life Cycle Inventory: Other Materials

- Example Latex coating material

	Proposed defaults	User entry
Quantity (kg/BoC, dry basis)	25.85	25.85
Moisture content as received (fraction between zero and one)	0	0
Upstream emissions for this raw material (kg CO <sub>2</sub> eq./kg dry)	2.628	2.628
Total received tonnes (wet)	0.02585	

	kg CO <sub>2</sub> eq./BoC		
	Scope 1	Scope 2	Scope 3
Upstream Emissions			57.60986
Truck, owned	0		0
Truck, non-owned			0.533979
Rail, owned	0		0
Rail, non-owned			0.079094
Water inland, owned	0		0
Water inland, non-owned			0
Water ocean, owned	0		0
Water ocean, non-owned			0
<b>Total</b>	<b>0</b>		<b>58.22294</b>
<b>Transport only</b>	<b>0</b>		<b>0.613073</b>

# Life Cycle Inventory: Electricity and Steam

- Need to know quantities and location of electricity

Quantity (MWh/BoC)	Region supplying the electricity	Default emission factor (kg CO <sub>2</sub> eq./MWh)		Selected emission factor (kg CO <sub>2</sub> eq./MWh)		
		Scope 2	Scope 3	Scope 2	Scope 3	Used for
0.0183	Alabama	711.0	18.2	711.0000	18.2000	Combined operations
0.0518	Kentucky	1045.4	25.3	1045.4000	25.3000	Combined operations
0.0157	Maryland	711.5	18.2	711.5000	18.2000	Combined operations
0.0306	Maine	393.6	11.5	393.6000	11.5000	Combined operations
0.0515	Michigan	738.2	18.8	738.2000	18.8000	Combined operations

kg CO <sub>2</sub> eq./BoC		
Scope 1	Scope 2	Scope 3
	11.03396573	0.282444692
	45.92225396	1.111376531
	9.472955502	0.242315938
	10.21380252	0.298421568
	32.23979045	0.821062125

# Life Cycle Inventory: Electricity and Steam

- For steam used a proxy:

Quantity (GJ/BoC)	Steam supplier/Source of emission factor	Default emission factor (kg CO <sub>2</sub> eq./MGJ)	Selected emission factor (kg CO <sub>2</sub> eq./GJ)
		Scope3	Scope 3
0.0434	<i>used natural gas EF</i>	<i>No default</i>	63.324



# Life Cycle Inventory: Manufacturing Wastes

- On site landfill that decays

	Proposed default	Selected value
Quantity of manufacturing wastes placed in industry landfills (dry kg/BoC)	43.09	83.50
Fraction of carbon in wastes	0.275	0.275
Fraction of carbon in wastes permanently stored	0.50	0.50
Fraction of wastes from owned operations	<i>No default</i>	1.00

## Results

	Scope 1	Scope 3
Mass of methane emitted from mill landfills (kg CO <sub>2</sub> eq./BoC)	172.21875	0
Mass of carbon permanently stored in mill landfills (kg CO <sub>2</sub> eq./BoC)	42.09791667	
Scope 1 Biogenic CO <sub>2</sub> emitted (kg CO <sub>2</sub> eq./BoC)	23.1538542	

# Life Cycle Inventory: Product Transport

- All transport steps involved, default emission data used

Product descriptor:	# T23 Code 293 USDOT 99,04 and USEPA 06 (printer to customer) 91%
Product transported:	Advertising material, commercial or trade catalogues, and similar printed products
Quantity (dry kg/BoC):	784.26
Moisture content:	0.08

Mode	Proposed defaults		Owned transportation		Non-owned transportation	
	Fraction of quantity transported	Distance, km	Fraction of quantity transported	Distance, km	Fraction of quantity transported	Distance, km
Truck	1	403.9			1	403.9
Rail	0	0				
Freshwater (inland) shipping	0	0				
Marine (ocean) shipping	0	0				

	kg CO <sub>2</sub> eq./BoC		
	Scope 1	Scope 2	Scope 3
Truck	0		32.19272
Rail	0		0
Marine (ocean) shipping	0		0
Inland (freshwater) shipping	0		0
<b>Total</b>	<b>0</b>		<b>32.19272</b>

# Life Cycle Inventory: Product Transport

- All transport steps involved, default emission data used

Transportation mode	(kg CO <sub>2</sub> / km*tonne)		
	Combustion	Precombustion	Total
Truck	0.0805	0.013	0.0935
Rail	0.0191	0.0031	0.0222
Marine (ocean)	0.0163	0.0022	0.0185
Inland (freshwater)	0.0288	0.0046	0.0334
Small truck (EOL)			1.26

# Life Cycle Inventory: End of Life: Carbon in Products

- How much carbon exists in products. Needed for end of life and carbon storage in products.
- Half life, number of years for the existing paper in use to half
- C permanently stored (in landfills)

Product	Carbon content (fraction)	Half-life (years)	Carbon permanently stored (fraction)
bleached kraft board	0.50	2.54	0.12
bleached kraft paper (packaging & industrial)	0.48	2.54	0.61
coated mechanical	0.50	2.54	0.85
coated woodfree	0.50	2.54	0.12
average containerboard	0.50	2.54	0.55
newsprint	0.46	2.54	0.85
recycled boxboard	0.50	2.54	0.55
recycled corrugating medium	0.50	2.54	0.55

# Life Cycle Inventory: End of Life

- Define the amount recycled
- Define the amount burned for energy and landfilled
- Built in data about landfill emissions

*The final product is probably used and disposed of in:*

U.S.

Disposition	Fractions	
	Proposed defaults	User Selection
Recycling	0.4210	0.388
Landfill	0.4696	0.498
Burning w/ energy recovery	0.1094	0.114

Transport distances, km	
Proposed defaults	User Selection
32.18	32.18
32.18	32.18
32.18	32.18

## Burning assumptions:

- GHG emissions are mainly N<sub>2</sub>O.

## Landfill assumptions:

- Landfills are assumed to be completely anaerobic.

- Fraction of gas transformed to methane:

50%

- Fraction of methane oxidized to CO<sub>2</sub> in landfill covers

10%

# Life Cycle Inventory: End of Life

Mass of product remaining in use after 100 years (kg/BoC)	31.58112712
Mass product landfilled (kg product/BoC)	413.46172757
Mass carbon landfilled (kg C/BoC)	133.54813801
Mass carbon permanently stored (kg C/BoC)	16.02577656
Mass if carbon transformed to gas (kg C/BoC)	117.52236145
Mass of carbon transformed into methane (kg C/BoC)	58.76118072
Mass of carbon in methane not oxidized in landfill covers (kg C/BoC)	52.88506265
Mass of carbon transformed into CO <sub>2</sub> (kg C/BoC)	5.87611807
Mass of carbon in methane burned for energy recovery (kg C/BoC)	23.26942757
Mass of methane emitted (kg CH <sub>4</sub> /BoC)	39.48751345
Landfill methane (kg CO <sub>2</sub> eq./BoC)	987.1878361
Burning GHGs (kg CO <sub>2</sub> eq./BoC)	0.946478653
Transport GHGs (kg CO <sub>2</sub> eq./BoC)	69.88853723
<b>Total EOL (scope 3) GHG emissions (kg CO<sub>2</sub> eq./BoC)</b>	<b>1058.022852</b>
<b>Carbon storage (kg CO<sub>2</sub> eq./BoC)</b>	<b>58.76118072</b>



# Life Cycle Inventory: Analysis

- Check for completeness, consistency, errors.....
- Interpret....

Basis of calculation (BoC, kg)	861.82556			
	Total	Scope 1	Scope 2	Scope 3
Total emissions, including transport (kg CO <sub>2</sub> eq./BoC):	3100	793.5	188.6	2118
Of which, total transport (includes all transport components):	161.2	0	0	161.2
Emissions from fuel used in manufacturing (including transport)	839.2	647.5	0	191.7
Emissions from purchased electricity and steam	383.5	0	188.6	194.9
Emissions from wood and fiber production (including transport)	312.9	0	0	312.9
Emissions from other raw materials (including transport)	298.1	0	0	298.1
Emissions from manufacturing wastes	146	146		0
Emissions from product transport	62.53	0	0	62.53
Emissions from end of life (including transport)	1058			1058
Total carbon storage changes (kg CO <sub>2</sub> eq./BoC)	138.3			
Changes in forest carbon (kg CO <sub>2</sub> eq./BoC)	0	Method used to estimate amount of product in use: <i>Weighted avg first order</i>		
Carbon in products in use (kg CO <sub>2</sub> eq./BoC)	37.4			
Carbon in landfills from products at end of life (kg CO <sub>2</sub> eq./BoC)	58.76			
Carbon in mill landfills from manufacturing wastes (kg CO <sub>2</sub> eq./BoC)	42.1			


# Life Cycle Inventory: Analysis

- Check for completeness, consistency, errors.....
- Interpret....






# Life Cycle Analysis of Paper: Carbon Footprint Results



# Life Cycle Analysis of Paper: Catalog System Boundary



# Full Life Cycle Analysis of Paper: Carbon Footprint Results

Table ES-6. LCIA Results – Catalog, Coated Freesheet

Impact category	Unit	Total (unit/catalog)	1- Fiber procurement	2- Coated freesheet production	3- Production of catalogs	4- Transport and use	5- End-of-life	Storage in use and landfill
Global Warming (GW)	kg CO <sub>2</sub> eq.	4.89E-01	5.4%	43.6%	15.7%	1.2%	37.7%	-3.4%
Acidification (AC)	H <sup>+</sup> moles eq.	1.67E-01	7.6%	67.4%	21.1%	1.1%	2.9%	N/A
Respiratory effects (RES)	kg PM <sub>2.5</sub> eq.	6.52E-04	3.5%	77.9%	15.6%	0.3%	2.6%	
Eutrophication (EU)	kg N eq.	8.85E-04	1.9%	19.0%	6.2%	0.2%	72.8%	
Ozone depletion (OD)	kg CFC-11 eq.	2.63E-08	6%	53%	31%	4%	7%	
Smog (SM)	kg NO <sub>x</sub> eq.	2.10E-03	7.7%	36.4%	48.7%	1.8%	5.3%	
Fossil fuel depletion (FF)	MJ surplus	3.94E-01	9.3%	52.4%	29.8%	2.6%	5.9%	

Results obtained using the *ecoinvent* database only (see Section 9.3.1.2 for more details)

National Council for Air and Stream Improvement, Inc. (NCASI). 2010. *Life cycle assessment of North American printing and writing paper products*. Unpublished Report. Research Triangle Park, NC: National Council for Air and Stream Improvement, Inc.

# Summary

- GHG concentrations are rising abruptly
- From a scientific viewpoint these are expected to increase radiative forcing and global warming
- A carbon footprint of a service is a method to gauge the net GWP
  - Includes emissions
  - Includes storage
- The carbon footprint is a partial life cycle analysis and as should not be considered in isolation
  - Often there is a tradeoff between carbon footprint and other environmental impacts that should be considered